

SLELO PRISM Partners Share These Goals:

PREVENTION

Prevent the introduction of invasive species into the SLELO PRISM region.

EARLY DETECTION & RAPID RESPONSE

Detect new and recent invaders and rapidly respond to eliminate all individuals within a specific area.

COOPERATION

Share resources, expertise, personnel, equipment and information.

INFORMATION MANAGEMENT

Collect, utilize, and share information regarding surveys, infestations, control methods, monitoring and research.

CONTROL

Control invasive species infestations by using best management practices, methods and techniques to include:

ERADICATION - Eliminate all individuals and the seed bank from an area.

CONTAINMENT - Reduce the spread of established infestations.

SUPPRESSION - Reduce the density but not necessarily the total infested area.

RESTORATION

Develop and implement effective restoration methods for areas that have been degraded by invasive species and where suppression or control has taken place.

EDUCATION / OUTREACH

Increase public awareness and understanding of invasive species issues through volunteer monitoring, citizen science and community outreach.

SLELO PRISM



FOR MORE INFORMATION

CONTACT:

St. Lawrence Eastern Lake Ontario
Partnership for Regional Invasive
Species Management

SLELO PRISM

Main Office

(315) 387-3600 x7725
c/o The Nature Conservancy

St. Lawrence County CCE
315-379-9192

Jefferson County CCE
315-788-8450

Lewis County SWCD
315-376-6122

Oneida County CCE
315-736-3394

Oswego County SWCD
315-592-9663

Or Visit Us Online At

www.sleloinvasives.org

Cover Photo:

https://en.wikipedia.org/wiki/Corbicula_fluminea.

Inside right column top identification photo:
www.sleloinvasives.org . Identification chart:

Center for Biodiversity and Conservation,
nyis.info.



SLELO PRISM

St. Lawrence Eastern Lake Ontario Partnership for Regional Invasive Species Management

What You Should Know About Asian Clam (*Corbicula fluminea*)



SLELO PRISM

*"Teaming up to stop the
spread of
invasive species"*

What are Asian Clam?

Asian clams (*Corbicula fluminea*) are native to the freshwaters of Eastern and Southern Asia. They were likely intentionally introduced to the west coast of North America in the 1930's as a food source for Chinese immigrants. Asian clams have been collected in running waters of central and western portions of New York State.

Asian clams are filter feeders and therefore feed on phytoplankton and other primary food sources in the aquatic ecosystem. Unlike zebra and quagga mussels, Asian clams have a pedal foot that allows them to feed on organic material and tiny organisms such as microbes, protists and meiofauna.

They have a high filtering capacity and population density and can disrupt aquatic food webs. In dense populations, Asian clams excrete significant amounts of inorganic nutrients, such as nitrogen, that stimulate the growth of algae and macrophytes. Furthermore, Asian clams can clog pipes of water treatment systems and power stations causing expensive damages.

Steps You Can Take to Stop the Spread of Asian Clam:

Prevention of the transport and sale of Asian clams is the most effective way to stop their spread.

Control/Management:

Physical Control:

Benthic barriers can deplete oxygen levels providing some control over small populations.

Mechanical Control:

Mollusks can be removed from piping by passing wads through pipes under pressure.

Chemical Control:

Molluscicides can be effective.

If you find Asian clam, please report your observation at iMapinvasives.org



Asian Clam Identification:

Description: Asian clam have a oval-triangular clam shape with a dorsal break (umbo) at the peak of the shell. The outside of the shell is olive or a yellowish-brown color with raised radial bands. The inner shell is typically white with a blueish tint to it and have finely serrated lateral “teeth” located between the valves at the umbo.



- Centrally located beak or umbo on shell
- Triangular or rounded triangular shell shape
- Many, coarse concentric rings on outside of shell
- 3 brown/purple colored radial colorbands (juveniles)
- 2 pair long, finely serrated lateral “teeth” per side on right valve; 1 pair per side on left valve
- 3 pseudocardinal “teeth” per valve.
- Interior of shell bluish white.
- Most similar to native fingernail clams.

Basic Identification characteristics for Asian clams. Graphic: Center for Biodiversity and Conservation