Our Goals

PREVENTION

Prevent the introduction of invasive species into the SLELO PRISM region.

EARLY DETECTION & RAPID RESPONSE

Detect new and recent invaders and rapidly respond to eliminate all individuals within a specific area.

COOPERATION

Share resources, expertise, personnel, equipment, and information.

INFORMATION MANAGEMENT

Collect, utilize, and share information regarding surveys, infestations, control methods, monitoring, and research.

CONTROL

Control invasive species infestations by using best management practices, methods and techniques to include:

- ERADICATION Eliminate all individuals and the seed bank from an area.
- CONTAINMENT Reduce the spread of established infestations.
- SUPPRESSION Reduce the density but not necessarily the total infested area.

RESTORATION

Develop and implement effective restoration methods for areas that have been degraded by invasive species and where suppression or control has taken place.

EDUCATION / OUTREACH

Increase public awareness and understanding of invasive species issues through volunteer monitoring, citizen science and community outreach.

INNOVATION

Explore technologies to enhance invasive species prevention and management initiatives.

SLELO PRISM

Hosted by The Nature Conservancy
315 387 3600

www.sleloinvasives.org

www.swallowwortcollaborative.org

PRISM Coordinator: Rob Williams (x7725)

rwilliams@tnc.org; Outreach Coordinator:

Megan Pistolese (x7724), megan.pistolese@tnc.org;

Aquatia Coordinator: Printers Program (x7730)

Aquatic Coordinator: Brittney Rogers (x7730), brittney.rogers@tnc.org; Terrestrial Coordinator:

Robert Smith (x7723), robert.l.smith@tnc.org

Explore, Ob serve, Report

Learn to recognize and report invasive species in our region.

For details contact:

megan.pistolese@tnc.org

Stay Connected

- 1. Email megan.pistolese@tnc.org
- 2. Type "join e-mail list" in subject space.
- 3. Hit send and receive seasonal e-newsletters and event updates.

Scan QR Code For More Resources









Cover Photo: USDA-APHIS http://www.hungrypests.com6.jpg. Sucker sprouts: Daniel Herms, the Ohio State University, bugwood.org, Canopy dieback/woodpecker damage photos: Geoff McVey, Forest Manger Limerick Forest Works, Brockville, Ontario. Ash Tree Identification Photos: David L. Roberts, Ph.D. Senior Academic Specialist, Michigan State University Extension, treedoctor.anr.msu.edu.



Emerald Ash Borer (Agrilus planipennis)





SLELO PRISM
Protecting Our Lands & Waters

What is an Emerald Ash Borer (EAB)?

This Asian beetle, (*Agrilus planipennis*) infests and kills North American ash tree species (*Fraxinus sp.*) including green, white, black and blue ash, and their cultivars. The larval stage of EAB feeds under the bark of ash trees, cutting off the flow of water and nutrients.

Visible Signs of EAB Infestation

- Sucker sprouts grow from base of tree
- Loss of leaves and branches







- Extreme wood pecker damage
- S-Shaped tunnels under bark





EAB Identification:

Adult EAB:

Metallic Green body, with coppery red abdomen under wings.



• Size: ½ inch wide and 1/8th inch long; small enough to fit on a penny.



- Adults may be present from **May-September**.
- They make 1/8" D-Shaped exit holes in bark which are often located towards the crown of the tree and hard to see.

EAB Larvae:

- Color: Creamy white
- Size: 1 inch-long "worms" with bell-shaped segments






* Larvae make **S-shaped tunnels under bark;** larvae themselves are hard to see.

Ash Tree Identification:

Branches/buds are arranged directly across from one another (opposite orientation)



Leaves are compound, containing 5-11 leaflets depending on tree species



Ash One leaf, 9 leaflets

Green Ash One leaf, 7 leaflets



Black Ash One leaf, 7 leaflets

White Ash top/bottom One leaf. 7

leaflets

> Bark has distinct diamond shaped ridges

