

# Priority Conservation Area Evaluation

---

## Oneida Lake & Three Mile Bay Wildlife Management Area

# Contents

Introduction .....	1
About this PCA – Oneida Lake – Three Mile Bay WMA .....	2
By the Numbers .....	3
Invasive Species Management Progress .....	3
Conservation Significance and Natural Features.....	4
Terrestrial Resilience.....	4
Carbon Benefits.....	5
Rare Native Species and Communities .....	6
Invasive Species Abundance and Management.....	7
Invasive Species Surveys.....	7
Approaching Species.....	8
Invasive Species Management.....	9
Pale swallow-wort.....	10
Partner Management Efforts.....	11
Recommendations & Future Work .....	11
References .....	i
Appendix A: Resilience Definitions.....	ii
Appendix B: Invasive Species Tier Ranking .....	iii
Appendix C: Oneida Lake - Three Mile Bay WMA Full Invasive Species Lists.....	iv
Appendix D: PCA Total Score Formula .....	vi


## Introduction

The St. Lawrence Eastern Lake Ontario Partnership for Regional Invasive Species Management (SLELO PRISM) serves as one of New York State's eight PRISMs and is hosted by The Nature Conservancy. The SLELO PRISM region encompasses 7,600 square miles and includes all or part of five counties (Map 1).

The mission of the SLELO PRISM is to protect native habitats, biodiversity, natural areas, parks and refuges, habitats, waterbodies, farmland, and open space by using a collaborative and integrated approach to invasive species management. The emphasis of these activities is on prevention, early detection, rapid response, ecological restoration, and education.

Due to the expansive nature of the PRISM and the high number of invasive species present in the region, SLELO PRISM partners focus invasive species management efforts on sites that are ecologically significant or have a high conservation value. In addition, sites that are seed banks, vectors, or that pose a proximity threat to high-value sites are all factors involved in determining site-based management on both public and private lands. The SLELO partnership has named these sites [Priority Conservation Areas \(PCAs\)](#). Specifically, PCA's are viewed as "sites that have ecological importance such as unique habitat, grassland, Alvar, wetland, dune, freshwater spawning area, fen, bog, etc. and are often host to a rare, threatened or endangered species."

Invasive species survey and management work at PCAs is further refined to areas where human activities or site conditions increase the probability of an invasive species being introduced and/or becoming established – known as **Highly Probable Areas (HPAs)**. Examples for HPAs in aquatic areas include boat launches, fishing access sites, and coves with shallow slow-



Map 1. Location of SLELO PRISM in New York State.

moving waters, while HPAs in terrestrial areas include trailheads, parking areas, and campgrounds.

The purpose of this priority conservation area evaluation is to summarize select measures of conservation significance within the PCA and to objectively assess SLELO's progress toward invasive species prevention, survey, and management goals. The evaluation includes an overview of a respective PCA's conservation significance; a summary of invasive species distribution and abundance, including a quantitative analysis of management progress; and recommendations for future work. Summaries of invasive species distribution and management progress are compiled with data collected by SLELO PRISM staff and contractors, supplemented by public data from the iMapInvasives database. Survey and management activities conducted by partners and not reported to the iMapInvasives database are not included in this assessment.

## About this PCA – Oneida Lake – Three Mile Bay WMA

Oneida Lake – Three Mile Bay Wildlife Management Area (WMA) PCA includes the aquatic habitat of Oneida Lake and the mixed habitat of the approximately 3,996-acre Three Mile Bay WMA, located in the Towns of West Monroe and Constantia, Oswego County. The NYSDEC managed WMA is a popular destination for wildlife dependent recreational activities such as hunting and fishing. The property is generally low elevation with little variation in topography. There is dominant muck-peat and acidic sandy soil that influence the vegetative community of the site (NYSDEC).

Oneida Lake is the largest waterbody located wholly within New York State with a surface area greater than 50,800 acres. It is generally shallow, with an average depth of only 22-feet and a maximum depth of 55-feet. It is approximately 21 miles in length with over 77 miles of shoreline. Oneida Lake is a popular sport fishery, often visited for its abundant population of walleye (NYSDEC).

Approximately 98% of the PCA is natural landcover (NLCD, 2019). The mean elevation of Oneida Lake - Three Mile Bay WMA PCA is 377 feet and most common geophysical settings are clay/silt (74%), calcareous loam (22%), and sand (4%).

Oneida Lake - Three Mile Bay WMA was nominated as a SLELO PCA in 2012 and includes approximately 11 terrestrial and 31 aquatic highly probable areas (Map 2). It was surveyed wholly or partially by SLELO staff/contractors in 2012, 2014, 2016, 2017, 2018, 2020, and 2022.




Map 2. Location of HPAs at Oneida Lake – Three Mile Bay WMA PCA (2022).

# BY THE NUMBERS

PCA STATS	SURVEY STATS	MANAGEMENT STATS
<p><b>54,790 +</b> ACRES IN THE PCA</p> <p><b>98%</b> NATURAL LANDCOVER</p> <p><b>96%</b> OF PCA IN RCN*</p> <p><b>317,269 mt</b> OF STORED CARBON</p> <p><b>22</b> KNOWN RARE SPECIES OR COMMUNITIES</p>	<p><b>42 HPAs</b> 11 TERRESTRIAL 31 AQUATIC</p> <p><b>21.4</b> AVERAGE ACRES SURVEYED PER VISIT</p> <p><b>50</b> KNOWN INVASIVE SPECIES (19 AQUATIC + 31 TERRESTRIAL)</p> <p><b>23</b> SLELO TIERED SPECIES (10 AQUATIC + 13 TERRESTRIAL)</p>	<p><b>1</b> TIER SPECIES UNDER MANAGEMENT</p> <p><b>16</b> TOTAL SLELO MANAGEMENT ACTIONS SINCE 2016</p> <p><b>0.55</b> AVERAGE ACRES MANAGED PER VISIT</p> <p><b>95%</b> AVG. REDUCTION IN EXTENT AT MANAGED SITES</p>

## INVASIVE SPECIES MANAGEMENT PROGRESS



**PCA SCORE** Score is based on the PCA's ecological attributes and current progress towards invasive species management goals. Primary drivers influencing the score at this PCA include:

**A+**

- (+) Above average terrestrial resilience
- (+) Terrestrial invasive species management objectives met

\* Current score subject to future discretionary restoration adjustment

## Conservation Significance and Natural Features

The following section provides an overview of select characteristics – such as terrestrial resilience, carbon storage, and rare species and communities – that lend to this PCA’s conservation value.

### Terrestrial Resilience

The Nature Conservancy has identified a network of lands with unique topographies, geologies, and other characteristics that can withstand the impacts of climate change. The **resilient and connected network (RCN)** identifies where plant and animal species have the best chance to adapt in a changing climate (Anderson et al., 2016). The attributes of climate resilient lands can be degraded by invasive plants and/or forest pests and pathogens. Multiple factors contribute to a location’s overall resilience, including:

- **Landscape Diversity** – microhabitats and climate gradients available within a given area. The persistence of species increases in areas with high landscape diversity.
- **Local Connectedness** – the number of barriers and degree of fragmentation within a given area. A permeable (or connected) landscape promotes resilience by facilitating species movements.

Approximately **96%** of the terrestrial environment within Oneida Lake - Three Mile Bay WMA is included in the resilient and connected land network (Figure 1). Oneida Lake - Three Mile Bay WMA PCA has **above average terrestrial resilience, above average local connectedness, and slightly above average landscape diversity**, indicating a slightly elevated capacity to maintain species diversity, movement, and ecological function as the climate changes (Figure 2). For background information on this data, see Appendix A.




Figure 2. Site resilience, local connectedness, and landscape diversity scores for Oneida Lake – Three Mile Bay WMA PCA. Scores are expressed as the standard deviation above or below the average score.




Figure 1. Percent of Oneida Lake – Three Mile Bay WMA located in The Nature Conservancy’s resilient and connected land network.

## Carbon Benefits

Climate change is driven, in part, by increased carbon dioxide in the atmosphere from human sources. Forests are an effective means to sequester (i.e. store) carbon. Through the process of photosynthesis, trees pull CO<sup>2</sup> from the air and bind it in their tissues as branches, roots, etc. Forests also sequester and store carbon through their soil. Conservation or improved management actions that aim to increase carbon storage and/or avoid carbon release are an important component of a natural climate solutions strategy. It's estimated that in the United States, conservation, restoration, and management could support sequestration of 21% of net annual emissions (Fargione et al., 2018). Unfortunately, tree damage or death caused by invasive forest pests or diseases can reduce sequestration and storage capacity. Based on an analysis of National Forest Inventory Plots, forests impacted by insect disturbances sequestered 69% less carbon than trees with no disturbance (Quirion et al., 2021). In addition, the presence of terrestrial invasive plants has been documented to reduce forest regeneration success, which can lead to long-term reductions in forest carbon storage (Magdalena & Katharina, 2020).

Models of forest and soil carbon data indicate Oneida Lake - Three Mile Bay WMA PCA stores an estimated 317,269 metric tonnes (mt) of carbon, including 275,337 mt forest carbon (Williams et al., 2021) and 41,932 mt of soil carbon (Guevara et al., 2020) (Figure 3).




Figure 3. Sources and quantities of stored carbon at Oneida Lake - Three Mile Bay WMA PCA.

## Rare Native Species and Communities

Approximately 42% of threatened or endangered species are at risk due to invasive species. Invasive species are generally considered one of the greatest causes of endangerment, second only to habitat loss (Pimental et al., 2005). Invasives may impact endangered species through direct predation, disease or competition for space and resources, and more (Duenas, et al., 2021).

Species and communities in New York are assigned a state rank by the NY Natural Heritage Program (NYNHP) to reflect their rarity. Conservation status ranks include:

- S1** – Typically 5 or fewer occurrences, very few remaining individuals, acres, or miles of stream, or some factor of its biology making it especially vulnerable in NY.
- S2** – Typically 6 to 20 occurrences, few remaining individuals, acres, or miles of stream, or factors demonstrably making it very vulnerable in New York State.
- S3** – Typically 21 to 100 occurrences, limited acreage, or miles of stream in New York State.
- S4** – Apparently secure in New York State
- S5** – Demonstrably secure in New York State

Surveys by NYNHP and other conservation partners have identified **19 rare species** and **3 rare communities** at Oneida Lake - Three Mile Bay WMA PCA (Table 1).

**Oneida Lake –  
Three Mile Bay WMA  
CONTAINS:**

3

**RARE  
COMMUNITIES**

19

**RARE  
SPECIES**

Table 1. Number of rare species and communities documented at Oneida Lake – Three Mile Bay WMA PCA.

State Rank	Functional Group	Count
Critically Imperiled (S1)	Fish	1
	Vascular Plants	7
Imperiled (S2)	Birds	2
	Fish	3
	Reptiles	1
	Vascular Plants	1
	Wetland/Aquatic Community	1
Vulnerable (S3)	Birds	4
	Wetland/Aquatic Community	2



# Invasive Species Abundance and Management

## Invasive Species Surveys

Oneida Lake - Three Mile Bay WMA PCA wholly or partially by SLELO staff/contractors in 2012, 2014, 2016, 2017, 2018, 2020, and 2022. Staff focus early detection surveys on tier-ranked species at approximately 11 terrestrial and 31 aquatic highly probable areas (HPAs). For more information on the ranking system, see Appendix B. Additional incidental invasive species observations throughout the PCA are submitted by community scientists and other practitioners. According to the iMapInvasives database, the first invasive species observation at Oneida Lake - Three Mile Bay WMA was reported in 1940. As of June 2023, 31 terrestrial and 19 aquatic invasive species are known to be present in Oneida Lake - Three Mile Bay WMA (Figures 4a). Twenty-seven are low to moderate impact species and not tier ranked in the SLELO PRISM. Eight known species are ranked as tier 3 and 14 are ranked as tier 4 (Figure 4b). The total number of known species at each HPA ranges from one to ten with an average of 3 species per HPA. The total number of SLELO tier species at each HPA ranges from one to six with an average of 2.3 species per HPA. For a full list of species known at the PCA, see Appendix C.




Figure 4. (a) cumulative number of aquatic and terrestrial invasive species documented at Oneida Lake - Three Mile Bay WMA PCA 2002-2021 (source: SLELO and iMapInvasives). (b) Distribution of known invasive species by SLELO tier classification.

## Approaching Species

The following species are not found within the PCA geography but are within five-miles of the PCA boundary and approaching the area. They should be considered for early detection efforts by staff and partners.

Table 2. Species approaching Oneida Lake - Three Mile Bay WMA PCA (iMapInvasives, 2023).

Scientific Name	Common Name	SLELO Tier	Observations in Buffer
<i>Lonicera maackii</i>	Amur Honeysuckle	4	1
<i>Adelges tsugae</i>	Hemlock Woolly Adelgid	3	3
<i>Microstegium vimineum</i>	Japanese Stiltgrass	3	1
<i>Centaurea spp.</i>	Knapweed (species unknown)	4	16
<i>Brachypodium sylvaticum</i>	Slender Falsebrome	1	3
<i>Vincetoxicum spp.</i>	Swallowwort (species unknown)	3	16
<i>Salix purpurea</i>	Basket Willow, Purpleosier Willow	NT	1
<i>Litylenchus crenatae mccannii</i>	Beech leaf disease nematode	NT	1
<i>Lotus corniculatus</i>	Bird's Foot Trefoil	NT	1
<i>Robinia pseudoacacia</i>	Black Locust	NT	2
<i>Cirsium vulgare</i>	Bull Thistle	NT	1
<i>Cirsium arvense</i>	Canada Thistle, Creeping Thistle	NT	4
<i>Epilobium hirsutum</i>	Codlins And Cream, Hairy Willow-herb	NT	2
<i>Valeriana officinalis</i>	Common Valerian	NT	1
<i>Hedera helix</i>	English Ivy	NT	3
<i>Craspedacusta sowerbyi</i>	Freshwater Jellyfish	NT	1
<i>Lonicera japonica</i>	Japanese Honeysuckle	NT	2
<i>Euphorbia virgata</i>	Leafy Spurge	NT	1
<i>Artemisia vulgaris var. vulgaris</i>	Mugwort	NT	1
<i>Lathyrus latifolius</i>	Perennial Pea, Sweet-pea Everlasting	NT	1
<i>Conium maculatum</i>	Poison-hemlock	NT	1
<i>Ligustrum spp.</i>	Privet (species unknown)	NT	5
<i>Viburnum sieboldii</i>	Siebold's Viburnum	NT	1
<i>Vicia cracca ssp. cracca</i>	Tufted Vetch	NT	5
<i>Morus alba</i>	White Mulberry	NT	1
<i>Wisteria spp.</i>	Wisteria (species unknown)	NT	1

## ***Invasive Species Management***

Known invasive species within the PCA are prioritized for management based on their current or future impacts and the availability of effective control measures and management resources. Because resources are limited, all known invasive species cannot be targeted for management. Some HPAs are not surveyed and/or managed annually. In addition, if a species is present in high abundance within the PCA, only a sub-set of HPAs may be prioritized for management.

One tier 3 species are targeted for management by SLELO staff and contractors at select HPAs within this PCA: pale swallow-wort (Table 3). The following section provides an overview of invasive species control efforts conducted by SLELO staff and contractors. Additional management actions conducted by partners or volunteers and reported to iMapInvasives are summarized under the sub-section “*Partner Management Efforts*”.

*Table 3. Simplified invasive species management plan overview for Oneida Lake - Three Mile Bay WMA PCA.*

<b>Management Goal(s):</b>	
Suppress known infestations of target invasive species where they occur at HPAs within Oneida Lake - Three Mile Bay WMA PCA to minimize their spread to uninvaded, interior portions of the property; protect rare, threatened, or endangered species and communities; maintain climate resilience; and promote the establishment and recovery of native species.	
<b>Monitoring Plan:</b>	
Outcome monitoring will be conducted annually by SLELO staff and/or contractors. Measurements of extent (acres) and invasive plant percent cover will be collected for each infestation to evaluate management progress.	
To validate current management goals and objectives, staff will conduct strategic monitoring of interior portions of the PCA at least every three-years to assess the extent of target invasive species outside HPAs. The location and size of all infestations will be recorded.	
<b>Restoration Needs:</b>	
As part of the annual monitoring process, SLELO staff will evaluate the need for active restoration at each management site. Restoration will be prioritized first for sites that have reached their management objective. Active restoration will utilize a selection of native species appropriate for the PCA and specific site.	
<b>Management Objective(s):</b>	
<b>Pale swallow-wort</b>	Reduce the extent of pale swallow-wort at HPAs by 90% by 2028

### Pale swallow-wort

Three pale swallow-wort infestations have been documented at the PCA since 2016. As of 2022, two infestations are under active management (Figure 6). The remaining site is unmanaged but is being evaluated for control.




Figure 6. Number of known knotweed infestations at Oneida Lake – Three Mile Bay WMA PCA from 2016-2022 by management status.

From 2016-2022, invasive species distribution and management data was collected using handheld GPS and manually transcribed to paper-based maps. Presence and treated area measurements observed from 2016-2019 (Figure 7) reflect this data collection technique and do not indicate unchanging conditions on the ground. Beginning in 2020, SLELO staff and contractors adopted a mobile GIS data collection system to allow for increased spatial mapping detail and accuracy. Based on the best available data, after seven years of chemical control, overall swallow-wort extent at HPAs reached a peak of 2.41 acres in 2020 and has been reduced by approximately 95% to 0.12 acres in 2022.




Figure 5. Changes in knotweed extent (presence area) and total area treated at all HPAs 2015-2022. Trendline represents two-year rolling average of presence area.

### Partner Management Efforts

The following sub-section provides an overview of management actions performed by partners and/or volunteers reported to iMapInvasives. Due to data reporting and sharing limitations, a quantitative assessment of management progress cannot be completed for each species. When possible, a summary of acres treated annually is provided. Trends charts are not provided for species managed sporadically.

- **Water chestnut (2.05 acres)**
- **Yellow iris (0.01 acres)**

### Recommendations & Future Work

- **Conduct strategic monitoring outside of HPAs to validate current invasive species management goals.** SLELO staff and/or contractors will explore opportunities to survey interior portions of the PCA to better understand the extent of target management species outside of HPAs. Management actions that aim to suppress or contain invasive species at HPAs could be undermined if target species are more widely distributed throughout the PCA.

## References

- Anderson, M.G., Barnett, A., Clark, M., Prince, J., Olivero Sheldon, A. and Vickery B. 2016. Resilient and Connected Landscapes for Terrestrial Conservation. The Nature Conservancy, Eastern Conservation Science, Eastern Regional Office. Boston, MA. [https://easterndivision.s3.amazonaws.com/Resilient\\_and\\_Connected\\_Landscapes\\_For\\_Terrestrial\\_Conservation.pdf](https://easterndivision.s3.amazonaws.com/Resilient_and_Connected_Landscapes_For_Terrestrial_Conservation.pdf)
- Dewitz, J., and U.S. Geological Survey, 2021, National Land Cover Database (NLCD) 2019 Products (ver. 2.0, June 2021): U.S. Geological Survey data release, <https://doi.org/10.5066/P9KZCM54>
- Fargione, J. E., Bassett, S., Boucher, T., Bridgham, S. D., Conant, R. T., Cook-Patton, S. C., et al. (2018). Natural climate solutions for the United States. *Sci. Adv.* 4:eaat1869. <https://www.science.org/doi/epdf/10.1126/sciadv.aat1869>
- Guevara, M., C.E. Arroyo-cruz, N. Brunsell, C.O. Cruz-gaistardo, G.M. Domke, J. Equihua, J. Etchevers, D.J. Hayes, T. Hengl, A. Ibelles, K. Johnson, B. de Jong, Z. Libohova, R. Llamas, L. Nave, J.L. Ornelas, F. Paz, R. Ressler, A. Schwartz, S. Wills, and R. Vargas. 2020. Soil Organic Carbon Estimates for 30-cm Depth, Mexico and Conterminous USA, 1991-2011. ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAC/1737>
- Langmaier, M., & Lapin, K. (2020). A Systematic Review of the Impact of Invasive Alien Plants on Forest Regeneration in European Temperate Forests. *Frontiers in Plant Science*, 11. <https://doi.org/10.3389/fpls.2020.524969>
- NYSDEC. (n.d.). Three Mile Bay Wildlife Management Area. Three Mile Bay Wildlife Management Area - NYS Dept. of Environmental Conservation. Retrieved March 18, 2022, from <https://www.dec.ny.gov/outdoor/70681.html>
- NYSDEC. (n.d.). Oneida Lake. Oneida Lake - NYS Dept. of Environmental Conservation. Retrieved March 18, 2022, from <https://www.dec.ny.gov/outdoor/41034.html>
- Pimentel, D., Zuniga, R., and D. Morrison. 2005. Update on the environmental and economic cost associated with alien-invasive species in the United States. *Ecological Economics*, 52. pp 273-288. doi: <https://doi.org/10.1016/j.ecolecon.2004.10.002>.
- Quirion B.R., Domke G.M., Walters B.F., Lovett G.M., Fargione J.E., Greenwood L, Serbesoff-King K., Randall J.M., Fei S. (2021). Insect and Disease Disturbances Correlate With Reduced Carbon Sequestration in Forests of the Contiguous United States. *Frontiers in Forests and Global Change*. 4. doi: <https://10.3389/ffgc.2021.716582>
- Williams, C.A., N. Hasler, H. Gu, and Y. Zhou. 2020. Forest Carbon Stocks and Fluxes from the NFCMS, Conterminous USA, 1990-2010. ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAC/1829>

## Appendix A: Resilience Definitions

**Resilience Score:** A site's Resilience Score estimates its capacity to maintain species diversity and ecological function as the climate changes. It was determined by evaluating and quantifying physical characteristics that foster resilience, particularly the site's landscape diversity and local connectedness. The score is calculated within ecoregions based on all cells of the same geophysical setting and is described on a relative basis as above or below the average. For example, cells of granite bedrock were compared with all other cells of granite bedrock, and coastal plain sands were compared with other coastal plain sands. Our goal was to identify the places most resilient to climate change for each geophysical setting within each ecoregion.

**Local Connectedness:** Refers to the degree of fragmentation and strength of barriers that create resistance to movement within a landscape. A highly connected landscape promotes resilience by allowing species to move through the landscape and find suitable microclimates where they can persist. In this study, we calculate local connectedness by measuring the amount and configuration of human-created barriers like major roads, development, energy infrastructure, and industrial farming and forestry land. Read the methods for your region:

**Landscape Diversity:** Refers to the microhabitats and climatic gradients available in the immediate neighborhood surrounding any 30-m cell of land. The persistence of species in an area increases in landscapes with a wide variety of microclimates created by the topography (topo-climates), elevation and hydrology. In this study, we measure microclimates by counting the variety of small-scale landforms, measuring elevation range, and evaluating the density and configuration of wetlands in a 100-acre neighborhood around every point on the landscape.

**Forest Carbon:** Estimates of 2010 forest carbon stock and components (aboveground, coarse woody debris, and soil/other) are from Williams et al. (2021b) following methods described for the Southeast US in Gu et al. (2019). To estimate carbon stock, attributes were determined for all forested 30-m pixels in the continental United States. A forest carbon cycle model trained to match Forest Inventory and Analysis (FIA) data was used to predict carbon stocks for 2010 based on site-level attributes of forest type group, years since disturbance, and site productivity class. Results were iterated backward in time to provide continuous, annual reporting of forest carbon dynamics for each pixel. Most prior studies lacked spatial detail on the age of forest stands that persisted in a forested condition during the satellite data era, but this study used remotely sensed biomass to estimate the stand age condition of these persisting, intact forests, distinguishing relatively young stands (e.g., 30 to 50 years old) from older stands.

**Soil Carbon:** Estimates of soil organic carbon (SOC) for 0-30 cm topsoil layer at 250-m resolution for the conterminous USA (CONUS) are from Oak Ridge Lab ([Guevara et al. 2020](#)). The estimates are for the period 1991-2010 and were derived using the USDA Rapid Carbon Assessment (RaCA), which used over 6000 field soil samples and multiple environmental variables representative of the soil-forming environment coupled with a machine learning approach (i.e., simulated annealing) and regression tree ensemble modeling for optimized SOC prediction. Across the continental US, nearly 31% of SOC was found in forests, 28% in croplands, and 35% in grasslands and shrublands respectively.

**Total Carbon:** Estimates for total carbon in the carbon calculator use Forest Carbon 2010 for all cells with forest cover and Soil Carbon 2010 for all cells with non-forest cover. To combine the two datasets, we resampled the SOC data to a 30-m resolution to align with our other data products, and then removed developed lands using the 2016 National Land Cover Dataset (NLCD). Please note that resampling to a higher 30-m resolution introduces false accuracy as the original SOC data was at a lower 250-m resolution.

For more information, visit:

<https://www.conservationgateway.org/ConservationByGeography/NorthAmerica/UnitedStates/edc/reportsdata/terrestrial/resilience/Pages/default.aspx>

## Appendix B: Invasive Species Tier Ranking

In 2017 the PRISM network, state agencies and the New York Natural Heritage Program (NYNHP) formalized official definitions for invasive species tiers based on an invasive species impacts and abundance. The ranking system included four primary tiers (Figure 10):

- **Tier 1 – Early Detection/Prevention:** Highly invasive species located in a buffer around region but not in region itself, targeted for early detection and prevention activities
- **Tier 2 – Eradication:** Highly invasive species with low abundance in the region, management goal of eradication
- **Tier 3 – Containment:** Highly invasive species with medium abundance in the region, management goal of containment
- **Tier 4 – Local Control:** Highly invasive species with great abundance in the region, management goal of local control

SLELO PRISM further prioritizes certain invasive species on the tiers list for management. These species are selected through nomination and agreement with PRISM partners. The SLELO tiered species list is a sub-selection of species ranked at the NYS scale.

		Difficulty of Eradication / Cost of Control Abundance (In PRISM plus Buffer)			
		None in PRISM	Low (Eradication/ Full containment may be feasible)	Medium (Strategic management to contain infestations and slow spread in PRISM)	High (Established/widespread in PRISM; only strategic localized management)
Impact (current and future)	Very High or High	<b>TIER 1</b> <i>Early Detection/Prevention</i> Highest level of early detection survey efforts. Should conduct delineation surveys and assign to appropriate Tier if detected. a) Inside buffer, but not in PRISM b) Outside PRISM and Buffer, but close (eastern North America) c) Far outside PRISM and buffer (not in east NA), but introduction pathway exists	<b>TIER 2</b> <i>Eradication</i> Highest level of early detection response efforts. High impact species with low enough abundance and suitable treatment method available to make eradication feasible within the PRISM. Need delineation surveys to determine extent.	<b>TIER 3</b> <i>Containment</i> Target strategic management to slow the spread, as likely too widespread for eradication, but many surrounding regions could be at risk if left unattended. For plants, use the IPMDAT. Possible eradication candidate only if adequate resources and effective control methods available.	<b>TIER 4</b> <i>Local Control</i> Eradication from PRISM not feasible; focus on localized management over time to contain, exclude, or suppress to protect high-priority resources like rare species or recreation assets. Be strategic when deciding if / where to control.
	Medium	<i>Evaluate (Medium Impact)</i> Further evaluate impacts and PRISM resources to see if the species should be assigned to one of the other lists. If this species could feasibly become high impact with climatic or other environmental changes, consider moving to the appropriate High Impact row based on abundance. If too little is known, consider moving to "Monitor".			
	Unknown	X	<b>TIER 5</b> <i>Monitor</i> Species that need more research, mapping, and monitoring to understand their invasiveness. This includes naturalized species and cultivated-only species that are known to be invasive in other regions but are not yet invasive here. Invasiveness may change with environmental or genetic changes. Should monitor populations on a regular basis to see if they are starting to become invasive and assign to appropriate Tier if invasive infestations detected.		

Figure 6. Invasive species tier table developed by the PRISMs, state agencies, and New York Natural Heritage Program.

Tier definitions were formalized in 2017 by the NYS invasive species network of PRISMs, state agencies, and NYNHP.

**Buffer:** An area chosen by the PRISM that surrounds the PRISM and takes in certain counties, states and provinces. Most PRISMs are using about 100 miles as the buffer.

**Impact:** Use the PRISM-specific invasiveness rankings if available, or use NYS ranks (see nyls.info for existing ranks). For species that are not ranked yet, or PRISM-specific adjustments of state ranks are deemed necessary, use expert opinion and document justification. Low-impact species not included since cannot justify spending resources to control these.

**Abundance:** This is left as a qualitative metric, since assigning standardized values to categories is not feasible due to the diversity of species dispersal strategies and data gaps.

This ranking system takes into account populations that have escaped into natural areas, but not intentionally (and legally) distributed individuals. For example, a landscape planting would not be counted.



## Appendix C: Oneida Lake - Three Mile Bay WMA Full Invasive Species Lists

<b>Terrestrial Species</b>	<b>SLELO Tier</b>	<b>Source</b>	<b>First Detected</b>
<i>Common Buckthorn</i>	4	iMap	2002
<i>Common mullein</i>	NT	iMap	2012
<i>Common reed grass, phragmites</i>	3	iMap	2012
<i>Garlic Mustard</i>	NT	iMap	2012
<i>Honeysuckle spp.</i>	4	iMap	2012
<i>Pale Swallowwort</i>	3	iMMA	2020
<i>Periwinkle</i>	NT	iMMA	2020
<i>Purple Loosestrife</i>	4	iMap	2014
<i>Wild Parsnip</i>	4	iMap	2018
<i>Yellow Iris</i>	4	iMap	2016
<i>Morrows honeysuckle</i>	4	iMap	2002
<i>Glossy Buckthorn</i>	4	iMap	2018
<i>Coltsfoot</i>	NT	iMap	2012
<i>Japanese Knotweed</i>	3	iMap	2014
<i>Goutweed</i>	NT	iMMA	2020
<i>Bittersweet Nightshade</i>	NT	iMMA	2020
<i>Orange Daylily</i>	NT	iMap	2012
<i>Multiflora Rose</i>	NT	iMap	2012
<i>Autumn Olive</i>	NT	iMap	2012
<i>Japanese Barberry</i>	NT	iMap	2016
<i>Emerald Ash Borer</i>	4	iMap	2018
<i>Tree-of-heaven, Ailanthus</i>	3	iMap	2018
<i>Norway Maple</i>	NT	iMap	2018
<i>Jumping Worms</i>	5	iMap	2020
<i>Bouncing Bet</i>	NT	iMap	2015
<i>Gold-moss</i>	NT	iMap	2015
<i>Greater Celandine</i>	NT	iMMA	2020
<i>Spongy Moth</i>	NT	iMap	2022
<i>Garden Stonecrop</i>	NT	iMMA	2020
<i>Creeping Jenny, Moneywort</i>	NT	iMMA	2020
<i>Elecampane</i>	NT	iMMA	2022

<b>Aquatic Species</b>	<b>SLELO Tier</b>	<b>Source</b>	<b>First Detected</b>
<i>Common Carp</i>	NT	iMap	1985
<i>Curly Pondweed</i>	4	iMap	1978
<i>Eurasian Water-milfoil</i>	4	iMap	2001
<i>European Frogbit; Common Frogbit</i>	4	iMap	2007
<i>Mud Bithynia</i>	NT	iMap	1966
<i>Water Chestnut</i>	3	iMap	1999

<i>Starry Stonewort</i>	3	iMap	2005
<i>Zebra Mussel</i>	4	iMap	1991
<i>Brittle Naiad</i>	NT	iMap	2012
<i>Rudd</i>	NT	iMap	1990
<i>Chinese Mysterysnail</i>	NT	iMap	1978
<i>European Stream Valvata</i>	NT	iMap	1990
<i>Bloody-red Shrimp</i>	NT	iMap	2009
<i>Quagga Mussel</i>	4	iMap	2008
<i>Rusty Crayfish</i>	3	iMap	2017
<i>Spiny Waterflea</i>	3	iMap	2019
<i>Round Goby</i>	4	iMap	2013
<i>Sea Lamprey</i>	NT	iMap	1940
<i>Banded Mysterysnail</i>	NT	iMap	1962

## Appendix D: PCA Total Score Formula

The total score for PCA's is calculated as follows:

$$\text{Total Score} = \text{Average}(\text{RCD} + \text{IS}) + \text{DRA}$$

- **RCD = Average(Resilience + Local Connectedness + Landscape Diversity Scores)**
  - Terrestrial resilience data from Anderson et al. (2016) – accessible [here](#).
- **IS = Average(Invasive Species Management Progress Scores)**
  - Invasive species management scores are calculated based on progress achieved toward established PCA and species-specific objectives. Objectives are set using extent or density-based metrics
    - Extent Based Metrics
      - *Example: Reduce net invaded area of all pale swallow-wort infestations at HPAs by 80% by 2025.*
        - “Score” is measured as progress achieved toward the set objective
          - If the objective is 80% and invaded area has currently been reduced to:
            - 55% → 55/80 = 0.69 (69% or D+)
            - 65% → 65/80 = 0.81 (81% or B-)
            - 75% → 75/80 = 0.94 (94% or A)
    - Density Based Metrics
      - *Ex: Reduce swallow-wort density at current HPAs to 5% or less by 2028.*
        - “Score” is measured as the amount change needed between current average invasive species density and goal
          - If the objective is 5% density or less, and current density is:
            - 76-100 – needs to move 4 cover classes = F
            - 51-75 – needs to move 3 cover classes = D
            - 26-50 – needs to move 2 cover classes = C
            - 5-25 – needs to move 1 cover class = B
            - <5% - at goal = A
- **DRA = Discretionary Restoration Adjustment**
  - A discretionary adjustment applied to the total score that reflects restoration progress and/or native species recovery at the PCA
    - Add (+) to Total Score
      - Based on visual field observation, the majority of management sites exhibit an increase in native/desirable vegetation richness or cover
    - No Adjustment
      - Based on visual field observation, the majority of management sites exhibit an increase in native/desirable vegetation richness or cover
    - Add (-) to Total Score
      - Based on visual field observation, the majority of management sites exhibit an increase in non-native or invasive vegetation richness and/or cover